



Comparative Efficacy of Selected Biopesticides with Chemicals against Shoot and Fruit Borer, *Leucinodes orbonalis* (Guenee) on Brinjal

**Potluri Nikhitha Chowdary^{a++*}
and Anoorag Rajnikant Tayde^{a#}**

^a Department of Entomology, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92337

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102958>

Original Research Article

Received: 17/05/2023

Accepted: 08/07/2023

Published: 15/07/2023

ABSTRACT

The present experiment was conducted during *Kharif* season of 2022 at Central Research Farm (CRF) of Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj. Total of two sprays were applied using eight treatments with three replications using Randomized Block Design (RBD), to evaluate the per cent infestation of shoot and fruit borer on brinjal. The results revealed that all the treatments were superior over the control against the infestation of brinjal shoot and fruit borer on third, seventh and fourteenth days after spraying. Among all the treatments, Chlorantraniliprole 18.5% SC @ 0.4ml/L was found to be the most effective treatment with (8.26%) infestation in shoot and (7.69%) infestation in fruit, followed by Emamectin benzoate 5% SG @

⁺⁺ M.Sc. Scholar;

[#] Assistant Professor;

*Corresponding author: E-mail: nikhithapraveen1229@gmail.com;

0.4gm/L (14.19%) and (12.55%). The next best treatments were found to be Indoxacarb 14.5% SC @ 1ml/L (14.39%) and (13.67%), Chlorpyrifos 20 EC @ 0.25ml/L (17.42%) and (14.40%), *Metarhizium anisopliae* @ 2×10^8 CFU (18.16%) and (14.64%), *Beauveria bassiana* @ 2×10^8 CFU (19.40%) and (15.46%). Neem oil 2% @ 20ml/L (21.88%) and (16.41%) was found to be least effective. The highest yield and cost benefit ratio was recorded in Chlorantraniliprole 18.5% SC (210.6 q/ha) and (1:7.48) followed by Emamectin benzoate 5% SG (187.5 q/ha) and (1:6.88), Indoxacarb 14.5% SC (163.5 q/ha) and (1:5.87), Chlorpyrifos 20% EC (131.9 q/ha) and (1:4.82), *Metarhizium anisopliae* (121.5 q/ha) and (1:4.28), *Beauveria bassiana* (116.3 q/ha) and (1:4.13) and Neem oil 2% (104.2 q/ha) and (1:2.67).

Keywords: Biopesticides; brinjal; chlorantraniliprole; cost benefit ratio; *Leucinodes orbonalis*.

1. INTRODUCTION

“Brinjal (*Solanum melongena* Linnaeus), also known as Eggplant, is referred as the “King of Vegetables”. It is originated in India and now grown as a vegetable throughout the tropical, sub-tropical and warm temperate areas of the world” [1]. The egg plant continues to be an important domestic crop cultivated across the country accounting for 9% of total vegetable production and covering 8.14% of land under vegetable cultivation.

“India is the second largest producer of brinjal in the world next to China” [2]. “The major brinjal growing states in India are West Bengal, Orissa, Gujarat, Bihar, Andhra Pradesh, Tamil nadu, Maharashtra and Madhya Pradesh. In India, West Bengal contributes the highest area 181.5 million hectare and production 2877 million tonnes, Karnataka has high productivity 25.4 million tonnes per hectare. In Uttar Pradesh, the area under cultivation of brinjal is 3430 hectare, producing 111.70 MT” [3].

“Though brinjal is a summer crop, it is been grown throughout the year under irrigated conditions. Hence it is subjected to attack by number of insect pest right from the nursery stage till harvesting” [4]. “Brinjal is attacked by more than 70 insect pests, among the insect pests infesting brinjal, the major one is shoot and fruit borer, *Leucinodes orbonalis* (Guenee)” [5]. “It is an internal borer and known to damage shoot and fruit of brinjal. The pest is estimated to cause 70 to 92 percent yield loss. The larvae of this pest cause 12 to 16 percent damage to shoots and 20 to 60 percent damage to fruits. The pest is very active during rainy and summer season and often causes more than up to 95 percent in India” [6].

“There is a tremendous misuse of insecticides in an attempt to produce damage free marketable

fruits. Insecticides have been reported effective against this pest but it is observed that this pest defies all the chemical control measures. Excessive dependence on huge quantities of insecticides, alone and in combinations, to control the pest is causing ecological pollution and pest resistance. It has become necessary to use preparations which are safe, effective and cheap” Sanjana and Tayde [7]. Hence present investigation was undertaken to evaluate the performance of certain insecticide at their recommended dosages against brinjal shoot and fruit borer.

2. MATERIALS AND METHODS

The present experimental work was carried out at Central Research Farm (CRF), Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj during *Kharif* season of 2022. The experiment was laid out in Randomized Block Design (RBD) with three replications. Crop was raised with a spacing of 60×45cm, in plots measuring 2m×1m each. The treatments *viz.*, Emamectin benzoate 5% SG @ 0.4gm/L, Indoxacarb 14.5% SC @ 1ml/L, Chlorpyrifos 20% EC @ 0.25ml/L, Chlorantraniliprole 18.5% SC @ 0.4ml/L, Neem oil 2% @ 20ml/L, *Metarhizium anisopliae* (2×10^8 CFU) @ 2.5 gm/L, *Beauveria bassiana* (2×10^8 CFU) @ 2.5 gm/L were applied twice at an interval of 15 days. The count of per cent infested shoots and fruits were recorded from five randomly selected and tagged plants. The observations were drawn on before and three, seven, fourteen days after spray. The per cent infestation on shoot and on fruit were calculated by using the following formulae,

On shoot: Shoot infestation was computed by counting the number of infested shoots and total number of shoots from five selected plants.

Table 1. Efficacy of different biopesticides and chemicals against brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guenee) (First spray: % shoot infestation) and (Second spray: % fruit infestation)

S. No	Treatments	Dose	First spray (% Shoot infestation)					Second spray (% Fruit infestation)					Yield (q/ha)	C:B ratio
			1DBS	3DAS	7DAS	14DAS	Mean	1DBS	3DAS	7DAS	14DAS	Mean		
T ₁	Indoxacarb 14.5% SC	1ml/L	25.93 (30.59)	15.36 ^c (23.04)	13.31 ^d (21.39)	14.50 ^b (22.38)	14.39 ^e	16.94 ^{de} (24.30)	14.96 ^{bc} (22.74)	12.42 ^{bc} (20.61)	13.63 ^{bc} (21.67)	13.67 ^{cd}	163.5	1:5.87
T ₂	Chlorpyrifos 20% EC	0.25ml/L	25.43 (30.26)	18.00 ^b (25.10)	17.05 ^c (24.30)	17.23 ^{cd} (24.45)	17.42 ^d	17.78 ^d (24.94)	15.62 ^{bc} (23.27)	13.01 ^{bc} (21.13)	14.59 ^{bc} (22.45)	14.40 ^{bcd}	131.9	1:4.82
T ₃	Chlorantraniliprole 18.5% SC	0.4ml/L	22.33 (28.17)	9.11 ^d (17.55)	7.06 ^e (15.23)	8.62 ^e (16.97)	8.26 ^f	13.96 ^f (21.94)	9.10 ^d (17.55)	6.92 ^d (15.95)	7.06 ^d (15.23)	7.69 ^e	210.6	1:7.48
T ₄	<i>Metarhizium anisopliae</i> 2×10 ⁸ CFU	2.5gm/L	26.49 (30.97)	19.31 ^b (26.06)	17.08 ^c (24.41)	18.11 ^c (25.18)	18.16 ^d	18.26 ^{cd} (25.29)	15.95 ^{bc} (23.52)	13.27 ^{bc} (21.24)	14.71 ^{bc} (22.51)	14.64 ^{bc}	121.5	1:4.45
T ₅	Neem oil 2%	20ml/L	24.66 (29.76)	23.39 ^a (28.89)	21.18 ^b (27.39)	22.07 ^b (28.00)	21.88 ^d	21.40 ^{ab} (27.54)	17.15 ^b (24.41)	15.70 ^b (23.24)	16.38 ^b (23.86)	16.41 ^b	104.2	1:3.76
T ₆	Emamectin benzoate 5% SG	0.4gm/L	22.48 (28.27)	15.20 ^c (22.91)	13.17 ^d (21.28)	14.20 ^d (22.13)	14.19 ^e	14.88 ^{ef} (22.69)	13.68 ^c (21.65)	11.00 ^c (19.29)	12.97 ^c (21.05)	12.55 ^d	187.5	1:6.88
T ₇	<i>Beauveria bassiana</i> 2×10 ⁸ CFU	2.5gm/L	28.77 (32.41)	20.18 ^b (26.69)	18.07 ^{bc} (25.15)	19.96 ^{bc} (26.51)	19.40 ^c	20.17 ^{bc} (26.66)	16.93 ^{bc} (24.27)	14.32 ^{bc} (22.19)	15.14 ^{bc} (22.90)	15.46 ^{bc}	116.3	1:4.26
T ₀	Control	-	23.51 (28.99)	25.30 ^a (30.20)	26.22 ^a (30.80)	27.16 ^a (31.40)	26.23 ^a	22.73 ^a (28.46)	25.05 ^a (30.03)	27.89 ^a (31.87)	29.23 ^a (32.72)	27.39 ^a	71	1:2.66
	F-test		NS	S	S	S	S	S	S	S	S	S	-	-
	S. Ed. (±)		2.09	1.17	1.45	1.64	0.62	1.00	1.51	1.89	1.40	1.14	-	-
	C. D. (P=0.05%)		-	2.51	3.12	3.52	1.09	2.16	3.25	4.06	3.00	2.00	-	-

*DBS=Day Before Spray, **DAS=Day After Spray, ***NS=Non-Significant, ****S=Significant

$$\% \text{ Shoot infestation} = \frac{\text{No.of shoots infested}}{\text{Total no.of shoots}} \times 100$$

(Rahman et al., [8])

On fruit: Fruit infestation was computed by counting the number of infested fruits and total number of fruits from five selected plants.

$$\% \text{ Fruit infestation} = \frac{\text{No.of fruits infested}}{\text{Total no.of fruits}} \times 100$$

(Rahman et al., [8])

The data on per cent infestation of brinjal shoot and fruit borer on both shoot and fruit were pooled separately and assessed to statistical analysis [9]. The fruit yield was harvested from each plot separately and the mean of marketable yield was recorded. Cost Benefit Ratio is also drawn by dividing the gross returns with the total cost of cultivation.

3. RESULTS AND DISCUSSION

The data on the pooled mean of per cent shoot infestation of brinjal shoot and fruit borer on 3rd, 7th and 14th day after the first spray revealed that, all the chemical treatments were significantly superior over control. Among all the treatments (Table 1), the lowest per cent shoot infestation was recorded in Chlorantraniliprole 18.5% SC (8.26%), Emamectin benzoate 5% SG (14.19%), Indoxacarb 14.5% SC (14.39), Chlorpyrifos 20% EC (17.42%), *Metarhizium anisopliae* (18.16%), *Beauveria bassiana* (19.40%) and Neem oil 2% (21.88%) was found to be least effective but significantly superior over the control (26.23%).

The data on the pooled mean (Table 1) of per cent fruit infestation of brinjal shoot and fruit borer on 3rd, 7th and 14th day after the second spray revealed that, the lowest per cent of fruit infestation was recorded in Chlorantraniliprole 18.5% SC (7.69%), followed by Emamectin benzoate 5% SG (12.55%), Indoxacarb 14.5% SC (13.67%), Chlorpyrifos 20% EC (14.40%), *Metarhizium anisopliae* (14.64%), *Beauveria bassiana* (15.46%) and Neem oil 2% (16.41%) was found to be the least effective but significantly superior over the control (27.39%).

Similar results of Chlorantraniliprole 18.5% SC, the most effective of above results, was reported by Tripura et al. [2]. The next best was Emamectin benzoate 5% SG, similar results

were supported by Patra et al. [10], followed by Indoxacarb 14.5% SC, which lined with the findings of Jagarlamudi and Kumar [11], Chlorpyrifos 20% EC, supported with the findings of Sanjana and Tayde [7], *Metarhizium anisopliae*, which lined with the findings of Sharma and Tayde [12], *Beauveria bassiana*, similar results were supported by Vyas and Tayde [13] and the results shown that the least effective was found to be Neem oil 2%, which lined with the findings of Chandar et al. [14].

The highest yield and cost benefit ratio were recorded in Chlorantraniliprole 18.5% SC (210.6 qt/ha) and (1:7.48), similar findings were supported by Vyas and Tayde [13], followed by Emamectin benzoate 5% SG (187.5 qt/ha) and (1:6.88), this result was supported by the findings of Patra et al. [10], Sharma and Tayde [12]. Followed by Indoxacarb 14.5 SC (165.4 qt/ha) and (1:5.87), this result was supported by the findings of Jagarlamudi and Kumar [11], followed by Chlorpyrifos 20% EC (131.9 qt/ha) and (1:4.82), this result is lined with the findings of Sanjana and Tayde [7], *Metarhizium anisopliae* (121.5 qt/ha) and (1:4.28) which is in the line with the findings of Sharma and Tayde [12], followed by *Beauveria bassiana* (116.3 qt/ha) and (1:4.13) which is in the line with the findings of Tripura et al. [2], followed by Neem oil 2% (104.2 qt/ha) and (1:3.79), these findings are in support with Chandar et al. [14].

4. CONCLUSION

The results revealed that Chlorantraniliprole 18.5% SC was the most effective treatment against brinjal shoot and fruit borer producing maximum yield and recording the highest cost benefit ratio compared to other treatments. While Emamectin benzoate 5% SG and Indoxacarb 14.5% SC have shown average results and the least effective chemicals were found to be Chlorpyrifos 20% EC, *Metarhizium anisopliae* and *Beauveria bassiana*. Botanical Neem oil 2% was found to be the least effective among all the treatments.

ACKNOWLEDGEMENT

The authors are thankful to Dr. Ashwani Kumar, Associate Professor and Head, Department of Entomology, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, for providing the facilities during the course of investigation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anwar S, Mari JM, Khanzada MA, Ullah F. Efficacy of insecticides against infestation of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee (Pyralidae:Lepidoptera) under field conditions. Journal of Entomology and Zoology Studies. 2015;3(3):292-295.
2. Chandar AS, Kumar A, Singh U, Kakade AA, Nawale JS, Narode MK. Efficacy of certain chemicals and biopesticides against brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guenee). Journal of Entomology and Zoology Studies. 2020;8(5):220-223.
3. Gomez KA, Gomez AA. Statistical procedures for agricultural research. A Wiley Interscience Publication. 1976;20-30.
4. Jagarlamudi SMR, Kumar A. Efficacy of certain chemicals with bio-pesticides against *Leucinodes orbonalis* (Guenee) on brinjal. Journal of Entomology and Zoology Studies. 2021; 9(1):771-777.
5. Kolhe PS, Kumar A, Tayde AR. Field efficacy of certain chemicals and neem products against shoot and fruit borer (*Leucinodes orbonalis* Guenee) on Brinjal (*Solanum melongena* Linnaeus) in Trans Yamuna Region of Allahabad. International Journal of Current Micro-Biology and Applied Plant Sciences. 2017;6(9):1320-1327.
6. Patra S, Thakur NSA, Firake DM. Evaluation of bio-pesticides and insecticides against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) in meghalaya of North-Eastern India. International Journal of Bio-resource and Stress Management. 2016;7(5):1032-1036.
7. Raghupathy A, Palanisamy S, Chandramohan N, Gunathilagaraj K. A guide on crop pests. Sooriya desk Top Publishers, Coimbatore. 1997;264.
8. Rahman MM, Islam KS, Jahan M, Uddin MA. Efficacy of some botanicals in controlling brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. Progressive Agriculture. 2009;20(1and2): 35-42.
9. Sanjana PSVV, Tayde AR. Field efficacy of some insecticides, neem oil and spinosad against shoot and fruit borer, *Leucinodes orbonalis* (Guenee) on brinjal. Journal of Entomology and Zoology Studies. 2019; 7(5):563-566.
10. Sharma JH, Tayde AR. Evaluation of bio-rational pesticides against brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guenee) at Allahabad Agro climatic Region. International Journal of Current Micro Biology and Applied Plant Sciences. 2017;6(6):2049-2054.
11. Shridhara M, Hanchinal SG, Sreenivas AG, Hosamani AC, Nidagundi JM. Evaluation of newer insecticides for the management of of Brinjal shoot and fruit borer, (*Leucinodes orbonalis* Guenee). International Journal of Current Micro Biology and Applied Plant Sciences. 2019;8(3): 2582-2592.
12. Tripura A, Chatterjee ML, Pande R, Patra S. Biorational management of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) in mid hills of Meghalaya. Journal of Entomology and Zoology Studies. 2017;5(4):41-45.
13. Vyas P, Tayde AR. Evaluation of chemical insecticides and biopesticides in management of shoot and fruit borer, *Leucinodes orbonalis* (Guenee) in brinjal *Solanum melongena* (L.). The Pharma Innovation Journal. 2022;11(10): 898-901
14. Yadav R, Lyall H, Kumar S, Sanap RK. Efficacy of certain botanical insecticides against shoot and fruit borer, *Leucinodes orbonalis* (Guenee) on brinjal (*Solanum melongena* L.). The Bioscan. 2015;10(2): 987-990.

© 2023 Chowdary and Tayde; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/102958>