



Effect of Micronutrients and Bioinoculants on Seed Yield and Seed Quality of Chilli (*Capsicum annuum* L.) under Kymore Plateau Zone

Pragya Singh^{a#*} and D. P. Sharma^{b++}

^a Department of Horticulture, RVSKVV, Gwalior (MP), India.

^b DES, JNKVV, Jabalpur (MP), India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i82005

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100476>

Original Research Article

Received: 18/03/2023

Accepted: 24/05/2023

Published: 03/06/2023

ABSTRACT

Chilli (*Capsicum annuum* L.) is one of the most extensively cultivated and economically important spice crops worldwide. Chilli is an economically important crop cultivated worldwide for its valuable fruits. The present study aimed to investigate the effect of micronutrients and bioinoculants on seed yield and quality of chilli. The experiment was laid out in Randomized Complete Block Design (RCBD-Factorial) with three replications and twenty treatments, including four micronutrients (M₁-M₅) and three bioinoculants (B₀-B₃). The interaction effect of micronutrients and bioinoculants was found to be significant in enhancing the chilli seed yield and quality. The treatment combination M₅B₃ (ZnSO₄ (0.2%) + *Trichoderma viride* (TV) + *Pseudomonas fluorescens* (PF) + *Azotobacter chroococcum* (AC) (2.5 kg/ha + 2.5 kg/ha + 5.0 kg/ha)) showed the highest improvement. Therefore, it can be inferred that a combination of micronutrients and bioinoculants is recommended for increasing the seed yield and quality of chilli.

Contractual / Guest Teacher;

++ Director;

*Corresponding author: E-mail: spragya455@gmail.com;

Keywords: Micronutrients; bioinoculants; sustainable agriculture; seed yield and quality.

1. INTRODUCTION

Chilli (*Capsicum annuum* L., 2n=24) is a significant vegetable, spice, and cash crop in India, belonging to the Solanaceae family. It is widely grown in both rainfed and irrigated environments throughout the country, with dried chilli production centered in the southern states. Madhya Pradesh alone produces 726.90 thousand tonnes of green chilli annually [1], cultivated in an area of 33.64 thousand ha, with major producing districts including Mandsaur, Kargone, Dhar, Shajapur, Ujjain, Chhindwara, Rajgarh, Khandwa, and Betul. Chilli is known for its pungency due to capsaicin present in the fruit's placenta and pericarp, which has diverse medicinal uses, and is rich in ascorbic acid [2]. Capsanthin, a pigment, is responsible for the red colour. Chilli may be grown in various soils, although well-drained loam soil is optimal. Chilli colour extracts are gaining popularity as a natural food colour substitute [3,4]. The chilli variety Arka Lohit developed as a pure line selection from IHR 324 (local collection), was released in 1990 at the AICRP (VC) workshop for national-level recognition. This variety exhibits tall plants with straight, smooth fruits that have pointed tip. The fruits undergo a color transformation from dark green to deep red and possess a high level of pungency. Arka Lohit adapts well to both irrigated and rainfed environments. Micronutrients play a crucial role in plant growth and the production of profitable crops, even though they are required in small quantities. Their application in various vegetables, including chillies, has been found to have positive effects on growth, seed yield, and quality [5,6].

Micronutrients play a crucial role in plant growth and development, and they can be provided to plants through foliar sprays for immediate nutrient availability. Essential micronutrients like iron, zinc, and boron are required in small quantities but have specialized physiological functions (growth and metabolism, etc.) in plants. Iron is essential for chlorophyll production and is particularly important for red chilli carotenoid synthesis, indirectly enhancing the quality of red chillies and paprika. Zinc is involved in auxin and protein synthesis and helps maintain membrane integrity. Additionally, zinc acts as a component of several enzymes; including dehydrogenase, aldolase, isomerases, proteinase, peptidase, and phosphohydrolase, further emphasizing its importance in plant metabolism [7]. Plant Growth

Promoting Rhizobacteria (PGPR) plays a vital role in promoting healthy growth and increasing the yield of various crops [8]. These beneficial bacteria actively colonize plant roots and enhance plant growth and yield through the production of phytohormones. Moreover, they protect plants from phytopathogenic microorganisms by producing siderophores, antibiotics, enzymes, and fungicidal compounds. Additionally, PGPR is known for its ability to perform asymbiotic N₂ fixation. Utilizing PGPR as biofertilizers and bioenhancers is an eco-friendly alternative to chemical fertilizers for various crop plants. In light of these benefits, this experiment was conducted to investigate the effect of micronutrients and bioinoculants on chilli seed yield and quality.

2. MATERIALS AND METHODS

The experiment entitled "Effect of Micronutrients and Bioinoculants on Seed Yield and Seed Quality of Chilli (*Capsicum annuum* L.) under Kymore Plateau Zone" was conducted during the Rabi season at the Horticulture Complex, Maharajpur, Department of Horticulture, JNKVV, Jabalpur (MP). A field experiment was conducted using a Randomized Complete Block Design (RCBD-Factorial) with various treatment combinations. The experiment included micronutrient treatments (M₁ – No micronutrient, M₂ - FeSO₄ (0.2%), M₃ - (CaNO₃)₂ (0.2%), M₄ - Borax (0.1%), M₅ - ZnSO₄ (0.2%)), bioinoculant treatments (B₀ - No bioinoculant, B₁ - TV (2.5 kg/ha), B₂ - TV + PF (2.5 kg/ha +2.5 kg/ha), B₃ - TV+PF+AC (2.5 kg/ha +2.5 kg/ha +5.0kg/ha)), and their interactions (M₁B₀-M₅B₃). Each treatment was replicated thrice to minimize experimental errors. Healthy 30 days old seedlings were transplanted at a spacing of 60 cm x 45 cm in the experimental plots. Foliar sprays of micronutrients were applied at three stages of plant growth: 60, 90, and 120 days after transplanting. In each plot, five randomly selected plants were tagged to collect biometric data on seed yield parameters, germination percentage, Seed vigour index-I (Standard germination (%) X Seedling length (cm)), and Seed vigour index-II (Standard germination (%) X Seedling dry weight (g)). The germination percentage of seeds was calculated by the guidelines of ISTA. Paper towels were used to hold the seeds, and 100 seeds from three replications were tested. The first and second germination counts were recorded on the fourth

and tenth days, respectively, and the percentage of germination was calculated based on normal seedlings. The seedling index was determined using two methods described by Abdul-Baki and Anderson [9]. The Seedling Vigour index was calculated by multiplying the germination percentage with either the seedling length (cm) or the seedling dry weight (g). Overall, the experimental setup and data collection procedures were conducted meticulously to evaluate the effects of micronutrients and bioinoculants on seed yield and seed quality of chilli. The data were analyzed statistically by the method of analysis of variance. The significance of various treatments was judged and suggested by Fisher [10] applying 'F' test.

3. RESULTS AND DISCUSSION

3.1 Seed Yield Parameters

The application of micronutrients and bioinoculants, both individually and in combination, had a significant impact on the characteristics of chilli seed yield.

3.1.1 100 seed weight (g)

The data in Table 1 shows that there was a highly significant effect on 100 seed weight. The maximum 100 seed weight was recorded in the application of micronutrients M_5 (2.81 g), while the lowest was observed in M_1 (1.63 g). Zinc plays a role in auxin and protein synthesis and contributes to seed formation and maturity. This finding is consistent with previous studies by [11,12]. Similarly, the application of bioinoculants also significantly influenced the 100 seed weight. B_3 (2.92 g) had the highest seed weight, while B_0 (2.00 g) had the lowest. *Trichoderma viride* and PGPR promote the production of growth substances such as auxins and cytokinins in plants, which positively affect processes like cell division and enlargement, resulting in bolder and heavier seeds. These findings are supported by [13,14]. Furthermore, when micronutrients and bioinoculants (*Trichoderma viride* and PGPR) were combined, their interaction had an impact on the 100 seed weight. The treatment combination M_5B_3 exhibited the highest seed weight (3.40 g), while the combination M_1B_0 had the lowest (1.42 g).

3.1.2 Number of seed per fruit

Under the micronutrient treatment, the highest number of seeds per fruit was recorded in M_5 (35.77), while the lowest count was observed in M_1 (25.41). The utilization of bioinoculants also

had a significant impact on the number of seeds per fruit, with the maximum count noted as B_3 (36.33) and the minimum count as B_0 (30.07). The interaction effects of micronutrients and bioinoculants, specifically *Trichoderma viride* and PGPR, showed a significant influence on the number of seeds per fruit. The maximum number of seeds per fruit was recorded in the M_5B_3 (38.88), whereas the minimum count was observed in the M_1B_0 (21.04). Zinc plays a vital role in seed development by aiding in auxin and protein synthesis. Additionally, the inoculation of plants with bioinoculants can enhance seed quality by improving nutrient uptake and stimulating the production of growth-promoting chemicals. These findings align with previous studies conducted by [11,12,13,15,16].

3.1.3 Seed yield per plant (g)

In Table 1, the results indicate that the application of micronutrients had a substantial impact on seed yield per plant. Among the treatments, the highest seed yield per plant was recorded in M_5 (3.99 g), while the lowest yield was observed in M_1 (2.77 g). Similarly, the application of bioinoculants also significantly influenced seed yield per plant, with the maximum yield recorded in B_3 (4.38 g) and the minimum yield in B_0 (3.07 g). The combined effects of micronutrients and bioinoculants, specifically *Trichoderma viride* and PGPR, found a significant influence on seed yield per plant. The maximum seed yield per plant was observed in the M_5B_3 treatment combination (4.85 g), while the minimum yield was found in the M_1B_0 combination (2.48 g).

3.1.4 Seed yield per hectare (q)

The data presented in Table 1 provides clear evidence of the significant impact of micronutrients on seed yield per hectare. Among the treatments, the highest seed yield per hectare was recorded in M_5 (0.136 q), while the lowest yield was observed in M_1 (0.094 q). Similarly, the application of bioinoculants also demonstrated a considerable influence on seed yield per hectare. Treatment B_3 exhibited the highest seed yield per hectare (0.146 q), while the lowest yield was recorded in B_0 (0.103 q). In the combined effects of micronutrients and bioinoculants, the maximum seed yield per hectare was achieved in the M_5B_3 (0.162 q), whereas the minimum yield was observed in the M_1B_0 (0.083 q). Zinc, an essential component of protein synthesis and nitrogen fixation, plays a crucial role in improving seed set and nutrient

uptake, thereby contributing to increased facilitate enhanced nutrient uptake. These seed yield. Bioinoculants, on the other hand, findings are consistent with previous studies produce growth-promoting chemicals and conducted by [12,13,16,17,18].

Table 1. Individual and interaction effect of different micronutrients and bioinoculants on seed yield and seed quality parameters

Treatments	Seed Yield Parameters				Seed Quality Parameters		
	100 seed weight (g)	No. of seed/ fruit	Seed yield /plant (g)	Seed yield/ hectare (q)	Germi nation %	Seed vigour index-I	Seed vigour index-II
Micronutrients							
M ₁ No micronutrient	1.63	25.41	2.77	0.094	62.53	345.38	179.10
M ₂ FeSO ₄ (0.2%)	2.52	34.04	3.64	0.125	71.95	766.63	252.87
M ₃ (CaNO ₃) ₂ (0.2%)	2.58	34.92	3.71	0.128	72.12	809.23	258.01
M ₄ Borax (0.1%)	2.65	35.12	3.81	0.132	73.41	875.49	264.05
M ₅ ZnSO ₄ (0.2%)	2.81	35.77	3.99	0.136	74.52	908.77	268.25
S.Em±	0.09	0.13	0.005	0.003	0.16	19.41	1.00
C.D.5% level	0.25	0.38	0.013	0.009	0.48	55.79	2.88
Bioinoculants							
B ₀ No bioinoculant	2.00	30.07	3.07	0.103	65.49	577.75	211.71
B ₁ TV (2.5 kg/ha)	2.30	32.17	3.20	0.112	70.21	671.57	237.57
B ₂ TV + PF (2.5 kg/ha +2.5 kg/ha)	2.54	33.65	3.69	0.13	73.06	786.15	255.87
B ₃ TV + PF+AC (2.5 kg/ha +2.5 kg/ha +5.0kg/ha)	2.92	36.33	4.38	0.146	74.87	928.94	272.69
S.Em±	0.08	0.12	0.004	0.003	0.15	17.36	0.89
C.D.5% level	0.22	0.34	0.012	0.008	0.43	49.89	2.87
Interaction							
M ₁ B ₀	1.42	21.04	2.48	0.083	60.83	284.75	142.40
M ₁ B ₁	1.57	25.03	2.50	0.083	62.81	326.33	169.01
M ₁ B ₂	1.72	26.33	2.95	0.103	62.84	360.75	194.45
M ₁ B ₃	1.82	29.25	3.14	0.105	63.64	409.69	210.55
M ₂ B ₀	1.98	31.21	3.16	0.106	66.39	558.89	221.07
M ₂ B ₁	2.38	33.45	3.27	0.115	69.56	725.82	251.60
M ₂ B ₂	2.63	34.98	3.59	0.127	74.84	802.82	263.04
M ₂ B ₃	3.07	36.52	4.55	0.152	76.99	979.00	275.78
M ₃ B ₀	2.10	32.20	3.20	0.107	66.56	602.54	224.73
M ₃ B ₁	2.42	33.74	3.36	0.119	69.99	752.19	253.25
M ₃ B ₂	2.67	35.31	3.65	0.133	74.87	833.86	273.07
M ₃ B ₃	3.12	38.45	4.61	0.154	77.07	1048.32	280.98
M ₄ B ₀	2.18	32.84	3.24	0.108	66.66	718.55	235.01
M ₄ B ₁	2.50	33.72	3.37	0.121	74.34	767.28	254.77
M ₄ B ₂	2.75	35.35	3.87	0.138	75.31	964.60	273.39
M ₄ B ₃	3.18	38.56	4.77	0.159	77.35	1051.53	293.05
M ₅ B ₀	2.30	33.07	3.26	0.111	67.01	723.99	235.32
M ₅ B ₁	2.60	34.89	3.48	0.124	74.34	786.21	259.20
M ₅ B ₂	2.95	36.25	4.37	0.146	77.45	968.70	275.39
M ₅ B ₃	3.40	38.88	4.85	0.162	79.29	1156.17	303.10
S.Em±	0.17	0.26	0.009	0.006	0.33	38.82	2.00
C.D.5% level	N.S.	0.77	0.027	N.S	0.97	111.57	5.75

TV=Trichoderma viride, PF=Pseudomonas fluorescense, AC=Azotobacterchroococcum

3.2 Seed Quality Parameter

The combined application of micronutrients and bioinoculants resulted in significant improvements in seed quality parameters, including germination percentage, seedling vigour index - I, and seedling vigour index - II.

3.2.1 Seed germination percentage

Seed germination percentage was significantly influenced by the foliar application of micronutrients. Treatment M_5 exhibited the highest seed germination percentage (74.52%), while the lowest percentage was recorded in treatment M_1 (62.53%). Similarly, the application of bioinoculants also had a notable effect on seed germination percentage, with the highest value observed in treatment B_3 (74.87%) and the lowest in treatment B_0 (65.49%). The combined effects of micronutrients and bioinoculants demonstrated a significant impact on seed germination percentage, with the highest percentage recorded in the M_5B_3 (79.29%) and the lowest in the M_1B_0 (60.83%). The inclusion of zinc sulphate in the micronutrient treatments increased metabolic activity and germination speed due to a higher concentration of metabolites, promoting the restart of embryonic growth during germination. Zinc, known for its involvement in protein synthesis, facilitated a better source-sink interaction, resulting in improved seed growth and increased germination. The enhanced production of hormones such as gibberellins may have activated specific enzymes that promoted early germination, potentially explaining these results. These findings are consistent with previous studies conducted by [16,19,20,21,22].

3.2.2 Seed vigour index-I

In the foliar application of micronutrients, the treatment M_5 recorded the highest value of seed vigour index-I (908.77), while the lowest seed vigour index-I was recorded in M_1 (345.38). Regarding bioinoculants, the maximum seed vigour index-I was observed in B_3 (928.94), whereas the minimum seed vigour index-I was recorded in B_0 (577.75). When considering the combined effects of micronutrients and bioinoculants, the treatment combination M_5B_3 exhibited the highest seed vigour index-I (1156.17), while the lowest value was observed in the treatment combination M_1B_0 (284.75).

3.2.3 Seed vigour index-II

In the foliar application of micronutrients, treatment M_5 (268.25) exhibited the highest value

of seed vigour index-II, while treatment M_1 had the lowest value (179.10). Regarding bioinoculants, the maximum seed vigour index-II was recorded in treatment B_3 (272.69), whereas the minimum seed vigour index-II was observed in B_0 (211.71). The interaction effects between micronutrients and bioinoculants were found to be significant. The highest seed vigour index-II was recorded in the treatment combination M_5B_3 (303.10), while the lowest value was observed in the combination M_1B_0 (142.40). Zinc plays a crucial role in protein synthesis and has been associated with a better source-sink interaction, resulting in improved seed development and a higher vigour index. Bioinoculants can significantly enhance seed germination, and the notable increase in seed vigour can be attributed to their increased auxin production. These findings are consistent with previous studies conducted by [12,16,21,23,22,24]. The above results reveal that the interaction effect of micronutrients and bioinoculants was the most effective treatment combination for increasing chilli seed yield and quality.

4. CONCLUSION

The application of micronutrients and bioinoculants in chilli cultivation offers a promising and sustainable approach to enhancing seed yield and quality. The findings highlight that this approach significantly improves microbial population and leads to substantial enhancements in seed yield and quality compared to the control group. The most effective treatment combination involves specific application rates of $ZnSO_4$, *Trichoderma viride*, *Pseudomonas fluorescens*, and *Azotobacter chroococcum*. This research provides valuable insights for farmers, researchers, and policymakers, emphasizing the importance of nutrient management and biological interventions in sustainable agriculture. However, further research is needed to optimize the application methods and rates, timing of these treatments for different chilli varieties and environmental conditions. Long-term field trials, scalability, mechanistic studies, and economic analysis are necessary to gain a deeper understanding of the underlying mechanisms, long-term effects, and economic feasibility of implementing these practical strategies for maximizing chilli seed yield and quality in commercial farming.

ACKNOWLEDGEMENTS

I would like to sincerely acknowledge the Dean, College of Agriculture, Jabalpur, for generously

providing the necessary facilities that were crucial for the successful completion of my research. I am extremely grateful and deeply honoured to express my heartfelt gratitude to Dr. D.P. Sharma, Joint Director of Extension Services at JNKVV, Jabalpur, for his invaluable guidance and insightful advice throughout this study. Their support and expertise have been instrumental in the achievement of this research endeavour.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. APEDA; 2022. Available:https://agriexchange.apeda.gov.in/India%20Production/India_Productions.aspx
2. Bal S, Sharangi AB, Upadhyay TK, Khan F, Pandey P, Siddiqui S, Saeed M, Lee, HJ, Yadav DK. Biomedical and antioxidant potentialities in Chilli: Perspectives and Way Forward. *Molecules*. 2022;27:6380.
3. Zhao Y, Chen ZY. Roles of spicy foods and their bioactive compounds in management of hypercholesterolemia. *Journal of Agricultural and Food Chemistry*. 2018;66:8662-8671.
4. Sann EE, Khin TN, Mar AA. Characterization of extracted natural colour from capsicum and its application in food product. *J. Myanmar Acad. Arts Sci*. 2020;18(1C):323-333.
5. Singh P, Sharma DP. Effect of micronutrients and bioinoculants (*Trichoderma viride* and PGPR) on yield parameters of chilli (*Capsicum annum* L.). *International Journal of Chemical Studies*. 2019a;7(4):773-775.
6. Singh P, Sharma DP. Response of *Trichoderma viride* and plant growth promoting rhizobacteria (PGPR) on growth and yield of chilli cv. Arka Lohit. *Journal of Pharmacognosy and Phytochemistry*. 2019b;8(4):1495-1497.
7. Mousavi SR. Zinc in crop production and interaction with phosphorus. *Australian Journal of Basic and Applied Sciences*. 2011;5(9):1503-150.
8. Backer R, Rokem JS, Ilangumaran, G, Lamont J, Praslickova D, Ricci E, Subramanian S, Smith DL. Plant growth-promoting rhizobacteria: context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. *Front. Plant Sci*. 2018;9:1473.
9. Abdul-Baki AA, Anderson JO. Vigour determination of soybean seed by multiple criteria. *Crop Science*. 1973;13:630-633.
10. Fisher RA. Comment on statistical problems in agricultural experimentation (with discussion). Supplemental to the *Journal of the royal Statistical Society Series B*. 1935;173:154-157.
11. Hussain SA, Mohammad S, Rao BVR. Response of chilli (*Capsicum annum* L.) to micronutrients. *Indian Journal of Agronomy*. 1989;34(1):117-118.
12. Kiran J, Vyakaranahal BS, Raikar SD, Ravikumar GH, Deshpande VK. Seed yield and quality of brinjal as influenced by crop nutrition. *Indian Journal of Agricultural Research*. 2010;44(1):1-7.
13. Yadegari M, Rahmani Asadi H. Evaluation of bean (*Phaseolous vulgaris*) seeds inoculation with *Rhizobium phaseoli* and plant growth promoting *Rhizobacteria* (PGPR) on yield and yield components. *African Journal of Agriculture Research*. 2010;5(9):792-799.
14. Yasari E, Patwardhan. Effect of Azotobacter and Azospirillum inoculations and chemical fertilizers on growth and productivity of canola. *Asian Journal of Plant Science*. 2007;6:77-82.
15. Basavarajeshwari CP, Ajjappalavara PS, Dhotre M. Effect of micronutrients on seed production in chilli. *Bhuvaneshwar, India*. 2010;305-310.
16. Deepika C. Effect of zinc and boron on growth, seed yield and quality of Radish (*Raphanus Sativus* L.) Cv. Arka Nishanth. *Current Agriculture Research*. 2015;3(1).
17. Naga SK, Swain SK, Raju B, Varma S. Influence of micronutrients application on growth and seed yield in tomato (*Lycopersicon esculentum* MILL.). *International Journal of Development Research*. 2013;3(11):191-195.
18. Ananthi M, Selvaraju P, Sundaralingam K. Evaluation of seed biopriming with biocontrol agents and biopesticides spraying on pests and its effect on seed yield and quality in chilli. *Journal of Entomology and Zoology Studies*. 2017;5(4):667-672.
19. Yoganand DK. Effect of mother plant nutrition and growth regulators on plant growth, seed yield and quality of bell

- pepper cv. California Wonder. M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Dharwad, Karnataka India. 2001;626.
20. Dileep Kumar Masuthi A, Vyakaranahal BS, Deshpandey VK. Influence of pelleting with micronutrients and botanical on growth, seed yield and quality of vegetable cowpea. *Karnataka Journal of Agriculture Science*. 2009;22(4):898-900.
21. Gholami A, Shamsavani S, Nezarat S. The effect of plant growth promoting rhizobacteria (PGPR) on germination, seedling growth and yield maize. *International Scholarly and Scientific Research and Innovation*. 2009;3(1):9-14.
22. Raman J. Response of *Azotobacter*, *Pseudomonas* and *Trichoderma* on growth of apple seedling. International Conference on biological and life science IPCBEE, IACSIT Press, Singapore; 2012.
23. Asaduzzaman M, Alam MJ, Islam MM. Effect of *Trichoderma* on seed germination and seedling parameters of chilli. *Journal of Science Foundation*. 2010;8(2):141-150.
24. Pal AK, Bara BM, Chaurasia AK. Influence of different micronutrient on seed viability and vigour parameters in Chilli (*Capsicum annum* L.) under storage condition. *International Journal Pure Application Bioscience*. 2018;6(5):377-383.

© 2023 Singh and Sharma; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/100476>